

## A Novel Synthetic Approach to Medium Ring Lactones via Dienetricarbonyliron Complexes

Ming-Liang Lai, Shun-Chi Chang, Chung-Chen Hwu, Ming-Chang P. Yeh\*

Department of Chemistry, National Taiwan Normal University, 88 Section 4, Ding-Jou Road, Taipei, Taiwan 117, ROC


**Abstract:** Reaction of lithium diisopropylamide (LDA) with ( $\eta^4$ -1,3-butadiene)Fe(CO)<sub>3</sub> complexes bearing carboester functionalized side chains at C-5 gives eight- to ten-membered lactones as the major products after electrophilic quenching. However, treatments of the complexes with LDA, under an atmosphere of carbon monoxide, furnish bridged bicyclo[5.2.1]decane and bicyclo[6.2.1]undecane lactone skeletons as the major products. Copyright © 1996 Elsevier Science Ltd

Medium-sized (eight- to twelve-membered) lactones and cyclic ethers constitute the characteristic molecular fragment of numerous natural products of biological interest, e.g. brevetoxin A,<sup>1</sup> laurencin,<sup>2</sup> laurenynne,<sup>3</sup> octalactin A,<sup>4</sup> octalactin B,<sup>4</sup> decan-9-olide,<sup>5</sup> and (*Z*)-dec-4-en-9-olide.<sup>5</sup> The most widely used method for the synthesis of medium-sized lactones involves intramolecular cyclization of a hydroxy and a carboxylic acid functionality (esterification),<sup>6</sup> Bayer-Villiger oxidation of seven-membered ring ketones,<sup>7</sup> Claisen rearrangement of a vinyl ketal to an eight-membered ring lactone,<sup>8</sup> or an atom transfer cyclization to eight- and nine-membered lactones.<sup>9</sup> However, only few reports on the preparation of medium-sized lactones promoted by transition metals. Of these reactions, however, formation of macrocyclic lactones has so far been restricted to the Stille coupling reaction.<sup>10</sup> We report here a novel method for the construction of medium-sized lactones via intramolecular cyclization of ( $\eta^4$ -1,3-diene)Fe(CO)<sub>3</sub> complexes bearing carboester functionalities at the terminal position of the diene ligand.

To prepare the starting carboester functionalized ( $\eta^4$ -1,3-diene)Fe(CO)<sub>3</sub> complexes, we adopted the well-known strategy developed by Birch and Pearson.<sup>11</sup> Treatment of cation **1** with zinc-copper reagent **2** [PivOCH<sub>2</sub>Cu(CN)ZnI], prepared from pivalate iodide and zinc followed by transmetallation with CuCN·2LiCl,<sup>12</sup> gave complex **3**. Reduction of **3** with diisobutylaluminum hydride (DIBAL) at 0 °C generated alcohol complex **4**. Acylation of **4** with acylchlorides (RCOCl, R = *i*-Pr or *n*-Pr) furnished ester complexes **5a** (88% from **1**) and **5b** (85% from **1**). Complex **6** with one more carbon unit on the side chain was synthesized in 69% yield from cation **1** and the zinc-copper reagent [BrZn(CN)CuCH<sub>2</sub>CO<sub>2</sub>Et] using the similar method.

Our intramolecular cyclization study began with complex **5a** (Scheme 1). Reaction of **5a** with 1.5 molar equiv of LDA in THF/HMPA (3:1) at -78 °C under nitrogen produced the eight-membered ring lactone **7a**

(8%) and the bridged bicyclic lactone **8** (20%) with an incorporated CO at the C-8 position after acid quenching. It is important to mention that lactones **7a** and **8** resulted from anti addition of the ester enolate at the internal position (C-2) of the diene ligand and no addition at C-3 was found. Unlike our previous reports, intramolecular nucleophilic additions of ester enolate to  $(\eta^4\text{-}1,3\text{-diene})\text{Fe}(\text{CO})_3$  complexes with 3 or 4 atom side chains occurred exclusively at the C-3 position of the diene ligand.<sup>13</sup> Thus, a longer side chain allows the ester enolate to attack at the internal position (C-2) of the diene ligand to generate the postulated homoallyl anionic intermediate **9**. Protonation of **9** generated iron-hydride species **10**. Reductive elimination of **10** gave **7a** (path a). However, intramolecular alkene insertion into the iron-hydride bond of **10**, followed by CO insertion and reductive elimination produced **8** (path b).<sup>14</sup> Thus, the addition was performed under an atmosphere of carbon monoxide (14 psi), which increased the yield of **8** to 39% as the only product isolated. The same reaction mixture could be quenched with molecular oxygen or iodomethane to furnish acid derivative **7b** (31%) and methyl ketone derivative **7c** (15%), respectively, as the only product in each case. None of the bridged bicyclic lactones, such as **8** was isolated.



However, under thermally controlled reaction conditions (25 °C), intramolecular cyclization of **5a** using 1.5 molar equiv of LDA generated nine-membered ring lactones **11** and **12** in a 3:4 ratio in 39% yield. Lactones **11** and **12** derived from addition of the ester enolate at the less hindered terminal position (C-1) of the diene ligands agrees closely with the formation of more stable  $(\eta^3\text{-allyl})\text{Fe}(\text{CO})_3$  anions from intermolecular additions of carbon nucleophiles to  $(\eta^4\text{-}1,3\text{-diene})\text{Fe}(\text{CO})_3$  complexes under thermally controlled reaction conditions.<sup>15</sup> Quenching the postulated allyl anion species **13** with acid afforded nine-membered lactones **11** and **12**. Under the same reaction conditions, intramolecular cyclization of complex **5b** gave the eight-membered ring lactone **14a** and the bridged bicyclic lactone **15** as a diastereomeric mixture in each case at -78 °C after acid quenching (**14a/15** = 2 : 1, 61% overall yield, entry 1, Table 1). The same reaction mixture was quenched with molecular oxygen to produce eight-membered ring lactone **14b**<sup>16</sup> as a mixture of diastereoisomers (*trans/cis* = 3:1, entry 2, Table 1). Moreover, intramolecular cyclization of **5b** under thermally controlled reaction conditions (25 °C) produced nine-membered ring lactones **16** and **17** in 2:1 ratio in 44% overall yield (entry 3, Table 1).

Increasing the tether length by one with complex **6** (entries 4–6, Table 1), also underwent intramolecular cyclization in the similar way to produce nine-membered ring lactone **18** as the major product (25%) and a trace amount of the bridged bicyclic lactone **19** at -78 °C (entry 4, Table 1). The yield of **19** (18%) could be increased by introducing an atmosphere of CO before acid quenching (entry 5, Table 1). As expected, intramolecular cyclization of **6** at 25 °C generated ten-membered ring lactone **20** in 8% yield (entry 6, Table 1).

Scheme 1




Table 1. Intramolecular cyclization and electrophilic quenching of complexes 5 and 6.

Entry	starting complexes	conditions	electrophiles	Product	yield
1	5b	$-78\text{ }^{\circ}\text{C}$	$\text{H}^+$	 14a ( $\text{R} = \text{H}$ )      15	61%
2	5b	$-78\text{ }^{\circ}\text{C}$	$\text{O}_2/\text{H}^+$	14b ( $\text{R} = \text{COOH}$ )	36%
3	5b	$25\text{ }^{\circ}\text{C}$	$\text{H}^+$	 16      17	44%
4	6	$-78\text{ }^{\circ}\text{C}$	$\text{H}^+$	 18      19 trace	25%
5	6	$-78\text{ }^{\circ}\text{C} / \text{CO}$	$\text{H}^+$	19	18%
6	6	$25\text{ }^{\circ}\text{C}$	$\text{H}^+$	 20	8%

**Acknowledgment:** This research was supported by a grant from the National Science Council of Republic of China (Grant No. NSC 84-2113-M-003-005).

#### Reference and Notes

1. Lin, Y. Y.; Risk, M.; Ray, S. M.; van Engen, D.; Clard, J.; Golik, J.; James, J. C.; Nakanishi, K. *J. Am. Chem. Soc.* **1981**, *103*, 6773.
2. (a) Irie, T.; Suzuki, M.; Masamune, T. *Tetrahedron* **1968**, *24*, 4193.  
(b) Irie, T.; Suzuki, M.; Masamune, T. *Tetrahedron Lett.* **1965**, *6*, 1091.
3. Falahaw, C. P.; King, T. J.; Imer, S.; Islimyeli, S.; Thomson, R. D. *Tetrahedron Lett.* **1980**, *21*, 4951.
4. Tapiolas, D. M.; Roman, M.; Fenical, W. *J. Am. Chem. Soc.* **1991**, *113*, 4682.
5. Moore, B. P.; Brown, W. V. *Aust. J. Chem.* **1976**, *29*, 1365.
6. Buszek, K. R.; Sato, N.; Jeong, Y. *J. Am. Chem. Soc.* **1994**, *116*, 5511.
7. (a) Carling, R. W.; Holmes, A. B. *J. Chem. Soc., Chem. Commun.* **1986**, 565.  
(b) Paquette, L. A.; Wang, T.-Z.; Sivik, M. R. *J. Am. Chem. Soc.* **1994**, *116*, 11323.
8. Carling, R. W.; Holmes, A. B. *J. Chem. Soc., Chem. Commun.* **1986**, 325.
9. Pirrung, F. O. H.; Hiemstra, H.; Speckamp, W. N.; Kaptein, B.; Schoemaker, H. E. *Synthesis* **1995**, 458.
10. (a) Kalivretenos, A.; Stille, J. K.; Hegedus, L. S. *J. Org. Chem.* **1991**, *56*, 2883.  
(b) Stille, J. K.; Su, H.; Hill, D. H.; Schneider, P.; Tanaka, M.; Morrison, D. L.; Hegedus, L. S. *Organometallics* **1991**, *10*, 1993.
11. (a) Birch, A. J.; Cross, P. E.; Lewis, J.; White, D. A.; Wild, S. B. *J. Chem. Soc. (A)*. **1968**, 332.  
(b) Pearson, A. J. *Acc. Chem. Res.* **1980**, *13*, 463.  
(c) Pearson, A. J.; Ong, C. W. *J. Am. Chem. Soc.* **1981**, *103*, 6686.
12. Knochel, P.; Chou, T. S.; Chen, H. G.; Yeh, M. C. P.; Rozema, M. J. *J. Org. Chem.* **1989**, *54*, 5202.
13. (a) Yeh, M. C. P.; Sheu, B. A.; Fu, H. W.; Tau, S. I.; Chuang, L. W. *J. Am. Chem. Soc.* **1993**, *115*, 5941.  
(b) Yeh, M. C. P.; Chuang, L. W.; Hwu, C. C.; Sheu, J. M.; Row, L. C. *Organometallics* **1995**, *14*, 3396.
14. Satisfactory spectral data (IR,  $^1\text{H}$  and  $^{13}\text{C}$ NMR, and HRMS) were obtained for all compounds.
15. (a) Semmelhack, M. F.; Herndon, J. W. *Organometallics* **1983**, *2*, 363.  
(b) Semmelhack, M. F.; Herndon, J. W.; Springer, J. P. *J. Am. Chem. Soc.* **1983**, *105*, 2497.  
(c) Semmelhack, M. F.; Le, H. T. M. *J. Am. Chem. Soc.* **1984**, *106*, 2715.  
(d) Semmelhack, M. F.; Le, H. T. M. *J. Am. Chem. Soc.* **1985**, *107*, 1455.  
(e) Yeh, M. C. P.; Kang, K. K.; Hwu, C. C. *J. Chin. Chem. Soc.* **1991**, *38*, 475.
16. Rigorous proof of the structures of **7b** and *trans* **14b** were accomplished by X-ray diffraction analysis.

(Received in Japan 28 May 1996; revised 1 July 1996; accepted 5 July 1996)