


# Substituent effects on the decomposition of chemiluminescent tricyclic aromatic dioxetanes

Chung-Wen Sun,<sup>a</sup> Shun-Chi Chen<sup>b</sup> and Tai-Shan Fang<sup>a\*</sup>

**ABSTRACT:** Three tricyclic 1,2-dioxetane derivatives, 1a, 2a and 3a were synthesized from their corresponding 1,4-dioxin acenaphthylene compounds, 1, 2 and 3, by reaction with singlet-oxygen (<sup>1</sup>O<sub>2</sub>) in dichloromethane. Evidence for the formation of the dioxetanes 1a, 2a and 3a is provided by the chemiluminescence (CL) that corresponds to the emission from the electronically excited diesters 1b\*, 2b\* and 3b\*, which are decomposed thermally from the dioxetanes 1a, 2a and 3a, respectively. The highly strained 1,2-dioxetane ring decomposes from a twisted geometry by simultaneous cleavages of the O–O and C–C bonds, producing the electronically excited diester that emits CL. It was observed that the CL from compound 2a is red-shifted relative to that of compounds 1a and 3a suggesting a higher degree of stabilization for the excited state by the electron-donating methoxy group. Also, a study of the solvent effect on fluorescence shows a significant red-shift in compound 2b, indicating a more polar excited state. The kinetics of the thermal decomposition of the 1,2-dioxetanes clearly demonstrate that the CL characteristics of compound 2a are quite different from those of compounds, 1a and 3a. These results are consistent with the proposed intramolecular chemically initiated electron exchange luminescence (CIEEL) mechanism which is triggered by the electron-donating group of compound 2a. Copyright © 2013 John Wiley & Sons, Ltd.



**Keywords:** chemiluminescence; tricyclic dioxetane; chemically initiated electron exchange luminescence

## Introduction

The phenomenon of firefly bioluminescence (1,2) has led to the theoretical (3,4) and practical (5–7) study of the chemiluminescence (CL) of 1,2-dioxetanes and has been of interest to chemists for a long time. High-energy 1,2-dioxetane molecules are the focus of many investigations because of their unique ability to decompose thermally into electronically excited carbonyl products. Although 1,2-dioxetanes have been postulated as reaction intermediates for over 100 years, charge transfer (CT)/electron transfer (ET) and electron back-transfer (EBT) have been suggested as a chemically initiated electron exchange luminescence (CIEEL) mechanism for CL, but the details of the mechanism have not yet been fully clarified (8). It is likely that the effect of structural changes will help to elucidate the still-debated mechanism of this reaction. In general [2 + 2] cycloaddition of singlet-oxygen with electron-rich alkenes is the most convenient method for the synthesis of 1,2-dioxetanes (9).

The key concept of the intramolecular CIEEL mechanism is that thermal decomposition of 1,2-dioxetane proceeds through a twisted diradical transition state that yields intramolecular radical ion intermediates. Subsequent exothermic EBT annihilation leads to an excited state and CL (10–15). For example, the intramolecular CIEEL mechanism of the dioxetane substituted with the electron-donating moiety [*p*-(dimethylamino)phenyl] is illustrated, where CT from the electron-donating moiety to

\* Correspondence to: Tai-Shan Fang, Department of Chemistry, National Taiwan Normal University, Taipei 116, Taiwan, Republic of China. E-mail: scchemts@ntnu.edu.tw

<sup>a</sup> Department of Chemistry, National Taiwan Normal University, Taipei 116, Taiwan, Republic of China

<sup>b</sup> Department of Chemical Engineering, Ming Chi University of Technology, Taishan, Taipei 243, Taiwan, Republic of China

the O–O bond of the dioxetane promotes decomposition of the dioxetane into the excited ester (see Scheme 1) (16).

The CL from ozonization and singlet-oxygen oxidation of the unsubstituted tricyclic 1,4-dioxin acenaphthylene compound **1** was reported several years ago from our laboratory (17). In continuation of this work, 5-methoxy-substituted 1,4-dioxin compound **2** and 5-chloro-substituted 1,4-dioxin compound **3** were synthesized to study the substituent effects on the decomposition of dioxetanes. We report herein the CL of the rigid tricyclic aromatic dioxetanes compounds, **1a**, **2a** and **3a**, with different functional groups (–H, –OCH<sub>3</sub> and –Cl) to assess the influence on the intramolecular CIEEL mechanism of 1,2-dioxetanes.

## Experimental

### Materials

**1,4-Dioxin acenaphthylene, compounds 1–3.** Acenaphthenequinone (ANQ) was obtained from Aldrich. 5-Nitro ANQ was synthesized from ANQ and sodium nitrate in concentrated sulfuric acid by refluxing the reaction mixture for 2 h. 5-Methoxy ANQ was synthesized from purified 5-nitro ANQ and potassium hydroxide in methanol by refluxing the reaction mixture for 2 h (18). 5-Chloro ANQ was synthesized from ANQ and *N*-chlorosuccinimide (NCS) in concentrated sulfuric acid by refluxing the reaction mixture for 2 h (Scheme 2A). Compounds **1–3** were synthesized by the photo-cycloaddition reaction of ANQ with tetramethylethylene (TME) in benzene (Scheme 2B) (19).

**Tricyclic 1,2-dioxetane compounds 1a–3a.** Singlet-oxygen oxidation of the 1,4-dioxin compounds **1–3** to produce the tricyclic 1,2-dioxetane compounds **1a–3a** was carried out by using the ozone-triphenyl phosphite procedure (5). The triphenyl phosphite ozonide, (PhO)<sub>3</sub>PO<sub>3</sub>, was prepared by passing extra pure oxygen through a commercial Fischer Model 501 ozonizer and bubbling the effluent into a solution of the phosphite in the range of 195–223 K (dry ice/acetone bath). The solution was saturated with ozone for ~2 h, and the temperature was kept within the range of 238–253 K for ~30 min to undergo singlet-oxygen oxidation of the 1,4-dioxin compounds. Dioxetane adducts **1a–3a** were isolated by ice-chilled *n*-hexane extraction from low-temperature synthesis of ozonized triphenyl phosphite [Scheme 2C (1)].

**Naphthalene diester compounds 1b–3b.** Two methods were utilized for the synthesis of the naphthalene diester compounds **1b–3b**. The first method involved the thermal decomposition of the compounds **1a–3a** to the naphthalene diester compounds **1b–3b** [Scheme 2C (1)]. The second procedure was the irradiation of compounds **1–3** in dichloromethane solution in the presence of the photosensitizer (methylene blue) under an oxygen atmosphere [Scheme 2C (2)]. The <sup>1</sup>H-NMR, IR and MS spectral data for the naphthalene diester compounds **1b–3b** are shown below.

Compound **1b**: <sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>): 7.99–7.95 (*d*, 2H, Ar–H, *J* = 8 Hz), 7.90–7.86 (*d*, 2H, Ar–H, *J* = 8 Hz), 7.56–7.48

(*t*, 2H, Ar–H, *J* = 8 Hz), 1.74 (*s*, 12H, 4CH<sub>3</sub>), FT-IR (KBr, cm<sup>-1</sup>): 66 2922, 1725, 1693, 1579, 1280, SIMS *m/z* 298 [M]. 67

Compound **2b**: <sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>): 8.46–8.42 (*d*, 1H, 68 Ar–H, *J* = 8 Hz), 7.95–7.91 (*d*, 1H, Ar–H, *J* = 8 Hz), 7.83–7.79 (*d*, 1H, Ar–H, *J* = 8 Hz), 7.52–7.44 (*t*, 1H, Ar–H, *J* = 8 Hz), 6.87–7.0 6.83 (*d*, 1H, Ar–H, *J* = 8 Hz), 4.03 (*s*, 3H, OCH<sub>3</sub>), 1.75 (*s*, 6H, 71 2CH<sub>3</sub>), 1.68 (*s*, 6H, 2CH<sub>3</sub>), FT-IR (KBr, cm<sup>-1</sup>): 2935, 1718, 1583, 72 1242, SIMS *m/z* 328 [M]. 73

Compound **3b**: <sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>): 8.51–8.47 (*d*, 1H, 74 Ar–H, *J* = 8 Hz), 7.93–7.89 (*d*, 1H, Ar–H, *J* = 8 Hz), 7.81–7.77 (*d*, 1H, Ar–H, *J* = 8 Hz), 7.67–7.60 (*m*, 2H, Ar–H), 1.74 (*s*, 6H, 76 2CH<sub>3</sub>), 1.70 (*s*, 6H, 2CH<sub>3</sub>), FT-IR (KBr, cm<sup>-1</sup>): 3005, 1734, 1687, 77 1570, 1283, SIMS *m/z* 332 [M]. 78

### Measurements


The absorption spectra were measured on a Hewlett-Packard diode array spectrophotometer. The steady-state emission spectra were obtained using a Cary Eclipsed Spectrofluorimeter equipped with a temperature controller. The absorption and emission experiments were performed with prepared solutions containing 1.0 × 10<sup>-4</sup> M of diesters in dichloromethane. For the temperature-dependent kinetic measurements, the temperature of the sample solution was controlled to within ±0.5 K using an electronically thermostating single cell and monitored with thermocouples attached to the cell.

## Results and discussion


The isolated dioxetane compounds **1a**, **2a** and **3a** were formed by reacting compounds **1**, **2** and **3**, respectively, with singlet-oxygen (<sup>1</sup>O<sub>2</sub>) in dichloromethane at 238 K and decomposed thermally into electronically excited diesters **1b\***, **2b\*** and **3b\***, which then fluoresced to their corresponding ground state configurations **1b**, **2b** and **3b**. Experimental evidence shows that the CL spectral peaks of compounds **1a–3a** are consistent with that of the photoluminescence spectral peaks of the photoexcited compounds **1b–3b**. Figure 1 shows the UV/Vis absorption and photoluminescence spectra of compounds **1b–3b**, and the CL spectra of compounds **1a–3a** in dichloromethane. Therein, the CL and photoluminescence peaking at λ<sub>max</sub> are normalized with the same relative intensity. It shows that the CL spectral peaks of the dioxetanes are consistent with those of the photoluminescence spectral peaks of the photoexcited diesters, respectively.

The CL peak of dioxetane compound **2a** at 430 nm shows an obvious red-shift in comparison with compounds **1a** (375 nm) and **3a** (378 nm). Presumably, the methoxy group of compound **2a** can donate electrons to the naphthalene ring and extend the conjugation of electrons, causing the obvious red-shift of CL.

Additional experimental evidence was obtained by examining the solvent effect on the photoluminescence of diesters **1b–3b**. Normalized fluorescence spectra of diesters **1b–3b** in different polar solvents are shown in Fig. 2. All the emissions from (π, π\*) excited states are seen in the three diesters, however,



Scheme 1.




Scheme 2.


the fluorescence spectrum of compound **2b** shows a noticeable dependence on the solvent polarity. By increasing solvent polarity from benzene to acetonitrile, the fluorescence peak of compound **2b** is red-shifted from 420 to 455 nm. Diesters **1b** and **3b** show emission peaks at 375 and 378 nm in benzene, and do not display a substantial shift with increasing solvent polarity. We can conclude, therefore,

that this long-wavelength fluorescence band of compound **2b** has a prominent characteristics CT which is stabilized in polar solvent.

From Figs 1 and 2, it can be concluded that the CL as well as the photoluminescence originate from the singlet excited state, which should be a  ${}^1(\pi^* \rightarrow \pi)$  state. In addition, the experimental results indicate that the electron-donating methoxy group




**Figure 1.** UV/Vis absorption, photoluminescence spectra of compounds **1b–3b**, and CL spectra of compounds **1a–3a** in dichloromethane.



**Figure 2.** Photoluminescence spectral peaks of compounds **1b–3b** in going from nonpolar to polar solvents.

present at the 5-position is able to stabilize the singlet excited state of the 1,2-dioxetane decomposition products. In the chemiexcitation processes, when both the O–O and C–C bonds are cleaved simultaneously, the electron-donating group (–OCH<sub>3</sub>) at the 5-position can interact through the  $\pi$  electrons of the naphthalene ring to stabilize the singlet excited state ( $^1(\pi^* \rightarrow \pi)$ ). Hence, the CL emission of this tricyclic aromatic dioxetane highly depends on substitution on the 5-position of the naphthalene structure or polarity of solvent system. These observations, together with the CL spectra are consistent with an intramolecular CIEEL decomposition initiated by CT from the electron-donating group of compound **2a**, followed by EBT to produce the excited diester. Compounds **1a** and **3a** indicate that the CIEEL mechanism is not operating prominently and they are decomposed thermally without involvement of CT or ET.

The rate constants for the decomposition of compounds **1a–3a** were obtained at 313–353 K by measuring the decay of the CL intensity. The CL decay of compounds **1a–3a** was monitored at the maximum value of the CL in dichloromethane (compound **1a**– $\lambda_{\text{CL,max}}$ , 375 nm; compound **2a**– $\lambda_{\text{CL,max}}$ , 430 nm; compound **3a**– $\lambda_{\text{CL,max}}$ , 378 nm). Figure 3 shows the time course



**Figure 3.** CL thermal decay of compound **2a** at different temperatures, monitored at 430 nm.

of the CL emission intensity at 430 nm upon thermal decomposition of compound **2a**, measured at different temperatures (313–353 K).

According to the transition state theory, the overall mechanism for the decomposition of the tricyclic 1,2-dioxetane compound can be presented as eqn (1).




where B = tricyclic 1,2-dioxetane compound,  $X^\ddagger$  = the activated complex and  $C^*$  = excited intramolecular naphthalene diester compound. The Eyring equation (eqn 2) can be obtained by transition state theory (20).

$$\ln \frac{k}{T} = \left( \ln \frac{R}{N_A h} + \frac{\Delta S^\ddagger}{R} \right) - \frac{\Delta H^\ddagger}{RT} \quad (2)$$

where  $N_A$  = Avogadro's constant and  $h$  = Planck's constant


The activation parameters of dioxetanes **1a–3a** were obtained from the temperature dependence of the CL decomposition rate constants. The CL activation parameters were obtained by plotting  $\ln(k/T)$  vs  $1/T$  in the Eyring plots, as shown in Fig. 4. The kinetics of the thermal decomposition of the 1,2-dioxetanes studied at 313–353 K reveal a significantly lower value for the activation enthalpy ( $\Delta H^\ddagger$ ) of **2a**.



**Figure 4.** Eyring plots of the CL kinetics of compounds **1a–3a**.

**Table 1.** Activation parameters and rates of decomposition for 1,2-dioxetanes 1a–3a

Compound	$\lambda_{\max}$ (nm)	kcal/mol	$\Delta H^\ddagger$ kcal/mol	$\Delta S^\ddagger$ cal/mol/K	$\Delta G^\ddagger$ (25°C) kcal/mol	$k_{25^\circ\text{C}}$ s <sup>-1</sup>	$\tau_{1/2}$ (25°C) h
1a (H)	375	(76.3)	20.3	-10.2	23.4	$4.54 \times 10^{-5}$	4.24
2a (OCH <sub>3</sub> )	430	(66.5)	13.1	-31.8	22.6	$17.1 \times 10^{-5}$	1.13
3a (Cl)	378	(75.7)	18.8	-15.8	23.5	$3.79 \times 10^{-5}$	5.08



Scheme 3.

activation enthalpies of the dioxetanes. The chemiexcitation process of the intramolecular CIEEL decay was rationalized and described as being mainly due to a particular form of electron-donating substituent at the 5-position of acenaphtho moiety. Further investigation of present and the other aspects of the intramolecular CIEEL mechanism are underway in our laboratory.

### Acknowledgements

Financial support from the National Science Council, MingChi University of Technology and National Taiwan Normal University, Taiwan, Republic of China, is gratefully acknowledged. Prof. LA Singer's (USC) comment is also highly appreciated.

### References

1. Fraga H. Firefly luminescence: a historical perspective and recent developments. *Photochem Photobiol Sci* 2008;7:146–58.
2. Nakatsu T, Ichiyama S, Hiratake J, Saldanha A, Kobashi N, Sakata K. Structural basis for the spectral difference in luciferase bioluminescence. *Nature Lett* 2006;440:372–6.
3. Vico LD, Lin Y-J, Krogh JW, Lindh R. Chemiluminescence of 1,2-dioxetane: reaction mechanism uncovered. *J Phys Chem A* 2007;111:8013–19.
4. Isoe H, Takano Y, Okumura M, Kuramitsu S, Yamaguchi K. Mechanistic insights in charge-transfer-induced luminescence of 1,2-dioxetanones with a substituent of low oxidation potential. *J Am Chem Soc* 2005;127:8667–79.
5. Adam W, Kazakov DV, Kazakov VP. Singlet-oxygen chemiluminescence in peroxide reactions. *Chem Rev* 2005;105:3371–86.
6. Matsumoto M. Advanced chemistry of dioxetane-based chemiluminescent substrates originating from bioluminescence. *J Photochem Photobiol C* 2004;5:27–53.
7. Schuster GB. Chemiluminescence of organic peroxides. conversion of ground-state reactants to excited-state products by the chemically initiated electron-exchange luminescence mechanism. *J Am Chem Soc* 1979;101:366–73.
8. Tanaka C, Tanaka J, Matsumoto M. An intramolecular charge/electron transfer chemiluminescence mechanism of oxidophenyl-substituted 1,2-dioxetane. *Phys Chem Chem Phys* 2011;13:16005–14.
9. Clennan EL, Pace A. Advances in singlet oxygen chemistry. *Tetrahedron* 2005;61:6665–91.
10. Hewage HS, Wallace KJ, Anslin EV. Novel chemiluminescent detection of chemical warfare simulant. *Chem Commun* 2007;3909–11.
11. Adam W, Bronstein I, Trofimov AV. Solvatochromic effects on the electron exchange chemiluminescence (CIEEL) of spiroadamantyl-substituted dioxetanes and the fluorescence of relevant oxyanions. *J Phys Chem A* 1998;102:5406–14.
12. Silva SM, Wagner K, Weiss D, Beckert R, Stevani CV, Baader WJ. Studies on the chemiexcitation step in peroxyoxalate chemiluminescence using steroid-substituted activators. *Luminescence* 2002;17:362–9.
13. Nery ALP, Ropke S, Catalani LH, Baader WJ. Fluoride-triggered decomposition of *m*-silyloxyphenyl-substituted dioxetanes by an intramolecular electron transfer (CIEEL) mechanism. *Tetrahedron Lett* 1999;40:2443–6.
14. Nery ALP, Weib D, Catalani LH, Baader WJ. Studies on the intramolecular electron transfer catalyzed thermolysis of 1,2-dioxetanes. *Tetrahedron* 2000;56:5317–27.

Table 1 shows the activation parameters and rates of decomposition for 1,2-dioxetanes 1a–3a. The  $\Delta H^\ddagger$  value of compound 2a is significantly lower than those values for compounds 1a and 3a. The rate constants obtained for compound 2a is ~4–5 times faster than that of 1a, and 3a at 298 K. The activation parameters of the unsubstituted compound 1a and the chloro-substituted compound 3a show similar thermal stability at room temperature, with a Gibbs' activation energy ( $\Delta G^\ddagger$ ) that differed by only ~0.1 kcal/mol. However, the methoxy-substituted compound 2a has a  $\Delta G^\ddagger$  value of 22.6 kcal/mol at 25°C corresponding to a  $\tau_{1/2}$  value of ~1.13 h, and shows a much lower thermal stability than compounds 1a and 3a. These results obtained upon intramolecular decomposition of dioxetanes clearly demonstrate that the CL characteristics of compound 2a are quite different from those of compounds 1a and 3a. The more negative activation entropy ( $\Delta S^\ddagger$ ) for compound 2a, which compensates for the surprisingly low value of  $\Delta H^\ddagger$ , can be understood by the need for a specific conformation for the CT from the methoxy group of the naphthalene ring to the O–O bond of the dioxetane, which promotes decomposition of the dioxetane into two radicals confined within a solvent cage. Subsequent EBT annihilation between the radicals releases enough energy to excite the naphthalene emitter to its singlet excited state. These results are consistent with the intramolecular CIEEL mechanism in which the electron-donating methoxy group promotes O–O bond cleavage by resonance interaction through the  $\pi$  electrons of the naphthalene moiety. The intramolecular CIEEL mechanism is triggered by the increasing electron-donating ability, with subsequent EBT annihilation leading to an excited state diester and luminescence to the ground state (Scheme 3).

### Conclusions

We have shown that the 1,2-dioxetane-based CL color is modulated by a dioxetane bearing a substituted-naphthalene group. The color of the CL emission from compound 2a is just different due to the different fluorescence spectrum of the thermodynamic decomposition product 2b, caused by the stabilization effect of the methoxy group. The change of substituent on the naphthalene emitter causes a change in the color of the dioxetane-based CL, and also affects the decomposition reaction

- 1 15. Ciscato LFML, Weiss D, Beckett R, Baader WJ. Fenchyl substituted  
2 1,2-dioxetanes as an alternative to adamantyl derivatives for  
3 bioanalytical applications. *J Photochem Photobiol A* 2011;218:41–7.  
4 16. Zaklika KA, Thayer AL, Schaap AP. Substituent effects on the decom-  
5 position of 1,2-dioxetanes. *J Am Chem Soc* 1978;100:4916–18.  
6 17. Fang T-S, Mei W-P. Ozonation and chemiluminescence of 8,8,9,9-  
7 tetramethyacenaphtho-[1,2-b]-1,4-dioxin: evidence for dual reaction  
8 pathways. *Tetrahedron Lett* 1987;28:329–32.  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
18. Wang L, Wang X, Cui J, Ren W, Meng N, Wang J, Qian X. Preparation  
of chiral *trans*-5-substituted-acenaphthene-1,2-diols by baker's  
yeast-mediated reduction of 5-substituted-acenaphthylene-1,2-  
diones. *Tetrahedron* 2010;21:825–30.  
19. Fang T-S, Singer LA. Wavelength dependent photoprocesses in  
acenaphthaquinone. *J Am Chem Soc* 1978;100:6276–78.  
20. Adam W, Zinner K. *Chemical and biological generation of excited  
states*. New York: Academic Press, 1982;153–89.
- 66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129

UNCORRECTED PROOF

# Author Query Form

---

**Journal: Luminescence**





**Article: bio\_2568**

Dear Author,

During the copyediting of your paper, the following queries arose. Please respond to these by annotating your proofs with the necessary changes/additions.

- If you intend to annotate your proof electronically, please refer to the E-annotation guidelines.
- If you intend to annotate your proof by means of hard-copy mark-up, please refer to the proof mark-up symbols guidelines. If manually writing corrections on your proof and returning it by fax, do not write too close to the edge of the paper. Please remember that illegible mark-ups may delay publication.

Whether you opt for hard-copy or electronic annotation of your proofs, we recommend that you provide additional clarification of answers to queries by entering your answers on the query sheet, in addition to the text mark-up.


Query No.	Query	Remark
Q1	AUTHOR: Please give city, state if USA, and country for suppliers at first mention in the text.	
Q2	AUTHOR: Please define IR.	
Q3	AUTHOR: Please define MS.	
Q4	AUTHOR: Please provide caption for scheme 1, 2 and 3.	

USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION


Required software to e-annotate PDFs: Adobe Acrobat Professional or Adobe Reader (version 7.0 or above). (Note that this document uses screenshots from Adobe Reader X)

The latest version of Acrobat Reader can be downloaded for free at: <http://get.adobe.com/uk/reader/>

Once you have Acrobat Reader open on your computer, click on the [Comment](#) tab at the right of the toolbar:



This will open up a panel down the right side of the document. The majority of tools you will use for annotating your proof will be in the [Annotations](#) section, pictured opposite. We've picked out some of these tools below:




**1. Replace (Ins) Tool – for replacing text.**



Strikes a line through text and opens up a text box where replacement text can be entered.

**How to use it**

- Highlight a word or sentence.
- Click on the [Replace \(Ins\)](#) icon in the Annotations section.
- Type the replacement text into the blue box that appears.




**2. Strikethrough (Del) Tool – for deleting text.**



Strikes a red line through text that is to be deleted.

**How to use it**

- Highlight a word or sentence.
- Click on the [Strikethrough \(Del\)](#) icon in the Annotations section.



**3. Add note to text Tool – for highlighting a section to be changed to bold or italic.**




Highlights text in yellow and opens up a text box where comments can be entered.

**How to use it**

- Highlight the relevant section of text.
- Click on the [Add note to text](#) icon in the Annotations section.
- Type instruction on what should be changed regarding the text into the yellow box that appears.

dynamic responses of mark ups  
ent with the **VAR** evidence




**4. Add sticky note Tool – for making notes at specific points in the text.**



Marks a point in the proof where a comment needs to be highlighted.

**How to use it**

- Click on the [Add sticky note](#) icon in the Annotations section.
- Click at the point in the proof where the comment should be inserted.
- Type the comment into the yellow box that appears.





USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION


**5. Attach File Tool – for inserting large amounts of text or replacement figures.**



Inserts an icon linking to the attached file in the appropriate place in the text.

**How to use it**

- Click on the [Attach File](#) icon in the Annotations section.
- Click on the proof to where you'd like the attached file to be linked.
- Select the file to be attached from your computer or network.
- Select the colour and type of icon that will appear in the proof. Click OK.



**6. Add stamp Tool – for approving a proof if no corrections are required.**




Inserts a selected stamp onto an appropriate place in the proof.

**How to use it**

- Click on the [Add stamp](#) icon in the Annotations section.
- Select the stamp you want to use. (The [Approved](#) stamp is usually available directly in the menu that appears).
- Click on the proof where you'd like the stamp to appear. (Where a proof is to be approved as it is, this would normally be on the first page).

of the business cycle, starting with the  
 on perfect competition, constant return  
 production. In this environment goods  
 extra profits and the number of firms  
 he number of firms is determined by the model. The New-Key  
 otaki (1987), has introduced product  
 general equilibrium models with nomin  
 ed and supply shocks. Most of this literat




**7. Drawing Markups Tools – for drawing shapes, lines and freeform annotations on proofs and commenting on these marks.**

Allows shapes, lines and freeform annotations to be drawn on proofs and for comment to be made on these marks..

**How to use it**

- Click on one of the shapes in the [Drawing Markups](#) section.
- Click on the proof at the relevant point and draw the selected shape with the cursor.
- To add a comment to the drawn shape, move the cursor over the shape until an arrowhead appears.
- Double click on the shape and type any text in the red box that appears.



For further information on how to annotate proofs, click on the [Help](#) menu to reveal a list of further options:

